[CO12/SQP006]

Time: 1 hour

Higher т Chemistry Paper I Specimen Question Paper NATIONAL QUALIFICATIONS

Check that the answer sheet provided is for Higher Chemistry Paper I.

Fill in the details required on the answer sheet.

Reference may be made to the Chemistry Higher and Advanced Higher Data Booklet (1999 edition). [This will be provided October/November 1998.]

Rough working, if required, should be done only on this question paper, or on the rough working sheet provided—**not** on the answer sheet.

Instructions for the completion of Part 1 and Part 2 are given on pages two and eight respectively.



PART 1

In questions 1 to 30 of this part of the paper, an answer is given by indicating the choice A, B, C or D by a stroke made in INK in the appropriate place in Part 1 of the answer sheet—see the sample question below.

For each question there is only ONE correct answer.

This part of the paper is worth 30 marks.

SAMPLE QUESTION

To show that the ink in a ball-pen consists of a mixture of dyes, the method of separation would be

- A fractional distillation
- B chromatography
- C fractional crystallisation
- D filtration.

The correct answer is **B**—chromatography. A heavy vertical line should be drawn joining the two dots in the appropriate box in the column headed **B** as shown in the example on the answer sheet.

If, after you have recorded your answer, you decide that you have made an error and wish to make a change, you should cancel the original answer and put a vertical stroke in the box you now consider to be correct. Thus, if you want to change an answer \mathbf{D} to an answer \mathbf{B} , your answer sheet would look like this:



If you want to change back to an answer which has already been scored out, you should enter a tick (\checkmark) to the RIGHT of the box of your choice, thus:



- **1.** In which of the following structures would the nails corrode before the roof itself?
 - A Zinc roof with iron nails
 - B Iron roof with copper nails
 - C Zinc roof with copper nails
 - D Copper roof with iron nails
- 2. Different isotopes of the same element have identical
 - A nuclei
 - B electron arrangements
 - C numbers of neutrons
 - D mass numbers.
- **3.** A mixture of magnesium bromide and magnesium sulphate is known to contain 3 mol of magnesium and 4 mol of bromide ions.

How many moles of sulphate ions are present?

- A 1
- B 2
- C 3
- D 4
- 4. The graph below shows the variation of concentration of a reactant with time as a reaction proceeds.



The average reaction rate, in moll⁻¹s⁻¹, during the first 20 s is

- A 0.0025
- B 0.0036
- C 0.0075
- D 0.0090.

5. The graph shows the distribution of kinetic energies of the molecules in a sample of gas.



Which graph would show the kinetic energies of the molecules when the sample is cooled by $10 \,^{\circ}\text{C}$?



- 6. A student found that 310 kJ of energy was released on burning 10 g of propan-1-ol, CH₃CH₂CH₂OH.
 From this experiment, what is the enthalpy of combustion, in kJ mol⁻¹, of propan-1-ol?
 - A -310
 - В -1296
 - C –1860
 - D -3100

7.	Silicon carbide can be used as	9. $N_2(g) + 2O_2(g) \rightarrow 2NO_2(g)$
	 A a lubricant B a tip for cutting/grinding tools C a substitute for pencil "lead" D an electrical conductor. 	How many litres of nitrogen dioxide gas could theoretically be obtained by sparking 4 litres of nitrogen gas with excess of oxygen gas?
		(All volumes are measured under the same conditions of temperature and pressure.)
		A 2
8.	What is the amount of oxygen atoms in 0.5 mol of carbon dioxide?	B 4
	A 0.25 mol	
	B 0.5 mol	
	C 1 mol	10 Which Calls Cills in source in source in the
	D 2 mol	10. Which of the following contains approximately 3×10^{23} molecules?
		A 17 g NH ₃
		В 36 g H ₂ O
		C 4 g CH ₄
		D 22 g CO ₂

11. Which reaction is an example of a reaction which takes place during reforming?

$$\textbf{A} \quad \textbf{CH}_3 - \textbf{CH}_2 - \textbf{CH}_2 - \textbf{CH}_2 - \textbf{CH}_3 \quad \rightarrow \quad \textbf{CH}_3 - \textbf{CH}_2 - \textbf{CH}_3 \quad + \quad \textbf{CH}_2 = \textbf{CH}_2$$

$$C CH_3 - CH_3 \rightarrow CH_2 = CH_2 + H_2$$

 $D \quad n \text{ CH}_2 = \text{CH}_2 \quad \rightarrow \quad - (\text{CH}_2 - \text{CH}_2)_n -$

12. A compound used in the synthesis of thermosetting plastics is:



The name of this compound is

- A methanol
- B methanal
- C methanoic acid
- D methanone.
- **13.** Which of the alcohols can be oxidised to give a ketone?
 - A 2-methylbutan-1-ol
 - B 2, 3-dimethylpentan-1-ol
 - C 3-methylbutan-2-ol
 - D 2-methylbutan-2-ol
- **14.** Which kind of reaction is used to produce an ester from a carboxylic acid and an alcohol?
 - A Addition
 - B Condensation
 - C Hydration
 - D Hydrolysis
- **15.** The dehydration of butan-2-ol can produce two isomeric alkenes, but-1-ene and but-2-ene.

Predict which alkanol can similarly produce, on dehydration, a pair of isomeric alkenes.

- A propan-2-ol
- B pentan-3-ol
- C hexan-3-ol
- D heptan-4-ol

- 16. Destroying ozone may have serious consequences.
 - Which statement about ozone is untrue?
 - A It absorbs ultraviolet radiation.
 - B It has the formula O_3 .
 - C It is classified as a CFC.
 - D It can react with halogenoalkanes.
- 17. A part of the formula for nylon is shown.

$$\begin{array}{cccc} H & H & O & O \\ | & | & || & || \\ -N - (CH_2)_6 - N - C - (CH_2)_4 - C - \end{array}$$

This polymer is classed as a

- A synthetic addition polymer
- B synthetic condensation polymer
- C natural condensation polymer
- D natural addition polymer.
- **18.** Polyester fibres and cured polyester are both very strong.

Which kinds of structure do their molecules have?

	Fibre	Cured resin
А	three-dimensional	three-dimensional
В	linear	linear
С	three-dimensional	linear
D	linear	three-dimensional

19. Part of a polyester chain is shown below.

Which compound, when added to the reactants during polymerisation, would stop the polyester chain from getting too long?

D

20. Which process is represented by the following equation?



21. Which of the following **must** contain nitrogen?

- A A protein
- B An oil
- C A polyester
- D A carbohydrate

- **22.** Which of the following is **not** a raw material in the chemical industry?
 - A Air
 - B Ethene
 - C Methane
 - D Water

23. The enthalpies of combustion of C(s), $H_2(g)$ and $C_4H_9OH(\ell)$, butan-1-ol, in kJ mol⁻¹, are as follows.

$$C(s) + O_2(g) \rightarrow CO_2(g) \qquad \Delta H = a$$
$$H_2(g) + \frac{1}{2}O_2(g) \rightarrow H_2O(\ell) \qquad \Delta H = b$$
$$C_4H_9OH(\ell) + 6O_2(g) \rightarrow 4CO_2(g) + 5H_2O(\ell)$$
$$\Delta H = c$$

The enthalpy of formation of butan-1-ol is given by the equation:

$$4C(s) + 5H_2(g) + \frac{1}{2}O_2(g) \rightarrow C_4H_9OH(\ell)$$

Which of the following can be used to calculate the enthalpy of formation of butan-1-ol?

- A 4a + 5b c
- B 2a + 10b c
- C c 4a 5b
- D = 2a + 5b + c
- 24. In which of the reactions will the equilibrium be unaffected by a change in pressure?
 - A $2NO(g) + O_2(g) \rightleftharpoons 2NO_2(g)$
 - B $2NO_2(g) \rightleftharpoons N_2O_4(g)$
 - C $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$
 - D N₂(g) + $3H_2(g) \rightleftharpoons 2NH_3(g)$
- 25.

 $\operatorname{Cl}_2(\mathbf{g}) + \operatorname{H}_2\operatorname{O}(\ell) \rightleftharpoons \operatorname{Cl}^-(\operatorname{aq}) + \operatorname{ClO}^-(\operatorname{aq}) + 2\operatorname{H}^+(\operatorname{aq})$

Which substance would move the above equilibrium to the right?

- A Hydrogen chloride
- B Sodium bromide
- C Chlorine
- D Hydrogen
- **26.** The following acid solutions all have a concentration of $0.1 \text{ mol } \Gamma^{-1}$.

In which solution would the concentration of $H^+(aq)$ ions be less than $0.1 \text{ mol } l^{-1}$?

- A Ethanoic acid
- B Hydrochloric acid
- C Nitric acid
- D Sulphuric acid

27. A fully dissociated acid is diluted by the addition of water.

Which of the following would increase with increasing dilution?

- A The pH value
- B The electrical conductivity
- C The rate of its reaction with chalk
- D The volume of alkali which it will neutralise
- **28.** During a redox process in acid solution, iodate ions, $IO_3^{-}(aq)$, are converted into iodine, $I_2(aq)$.

$$2 \text{ IO}_3(\text{aq}) \rightarrow \text{I}_2(\text{aq})$$

The numbers of $H^+(aq)$ and $H_2O(\ell)$ required to balance the ion-electron equation for the formation of 1 mol of $I_2(aq)$ are, respectively

- A 6 and 3
- B 3 and 6
- $C \quad 12 \text{ and } 6$
- D 6 and 12.
- 29. Which of the following has an electrical charge?
 - A α -particles
 - B X-rays
 - C Neutrons
 - D γ-rays
- 30. The chart below was obtained from an 8-day old sample of an α -emitting radioisotope.



What is the half-life of the radioisotope?

A	2 days
B	4 days
С	8 days
D	12 davs

Page seven

PART 2

In questions 31 to 34 of this part of the paper, an answer is given by circling the appropriate letter (or letters) in the answer grids provided on Part 2 of the answer sheet.

In some questions, two letters are required for full marks.

If more than the correct number of answers is given, marks may be deducted.

In some cases the number of correct responses is NOT identified in the question.

This part of the paper is worth 10 marks.

SAMPLE QUESTION

A CH ₄	Н ₂	C CO ₂
D CO	E C ₂ H ₆	F N ₂

(a) Identify the diatomic **compound(s)**.

А	В	С
D	Е	F

The one correct answer to part (a) is D. This should be circled.

(b) Identify the two substances which burn to produce both carbon dioxide and water.

A	В	С
D	E	F

As indicated in this question, there are **two** correct answers to part (*b*). These are A and E. Both answers are circled.

(c) Identify the substance(s) which can **not** be used as a fuel.

А	В	C
D	Е	F

There are **two** correct answers to part (*c*). These are C and F.

Both answers are circled.

If, after you have recorded your answer, you decide that you have made an error and wish to make a change, you should cancel the original answer and circle the answer you now consider to be correct. Thus, in part (a), if you want to change an answer **D** to an answer **A**, your answer sheet would look like this:

A	В	С
Ø	Е	F

If you want to change back to an answer which has already been scored out, you should enter a tick (\checkmark) in the box of the answer of your choice, thus:

X	В	С
	Е	F



Curve I shows the results for the reaction of excess zinc with 100 cm^3 of 0.2 mol 1^{-1} sulphuric acid.

A		В		С	
	Excess copper		Excess magnesium		Excess iron
D		Е		F	
	$50 \text{ cm}^3 \text{ of } 0.2 \text{ mol I}^{-1}$ sulphuric acid		$100 \text{ cm}^3 \text{ of } 0.2 \text{ mol } 1^{-1}$ sulphuric acid		$200 \text{ cm}^3 \text{ of } 0.2 \text{ mol I}^{-1}$ sulphuric acid

(a) Identify the two chemicals which would react to give the results shown by curve II.

(b) Identify the two chemicals which would react to give the results shown by curve III.

32. The first twenty elements in the Periodic Table can be categorised according to their bonding and structure.



(a) Identify the element which exists as a covalent network solid.

(b) Identify the element which exists as a discrete covalent molecular solid.

(c) Identify the most electronegative element.



- (a) Identify the compound which is an isomer of the compound shown in box A.
- (b) Identify the compound(s) which could be oxidised to form the compound shown in box **E**.
- 34. Read the following passage from a popular scientific journal. It discusses the nuclear reactions which occur during the explosion of a star.

"Stars produce heavier elements when they explode as supernovae . . . The main product (of the supernova) should be radioactive nickel. It forms (from) nuclei of oxygen. Heat from a shock wave 'welds' the oxygen into heavier nuclei. This radioactive nickel decays into cobalt, which in turn decays into iron."

(from New Scientist, 11 August 1988)

Identify the reaction(s) which can be found in the passage.

А	Nuclear fission	В	Nuclear fusion
С	Loss of an alpha particle	D	Loss of a beta particle
Е	Loss of a proton	F	Loss of a neutron

[END OF QUESTION PAPER]

NATIONAL

QUALIFICATIONS

Higher Chemistry Paper 1

Specimen Question Paper

Ð

ANSWER SHEET

Full name of school or college	Town
First name and initials	Sumame
Date of birth	Number of seat
Using ink, indicate your choice of answer A by a single stroke joining the two dots in the box, as in the following example:	B C D Using ink, indicate your choice of answer(s) by circling the appropriate letter(s) as in the following example:
A B C D A E	B C D A B C
	· · · З1а А В С ЗЗа А В С
	b A B C
	C A B C
15	

[CO12/SQP006]

Higher	Time: 1 hour 30 minutes	NATIONAL
Chemistry		QUALIFICATIONS
Paper II		
Specimen Questio	on Paper	

Fill in these boxes and read what is printed below.			
Full name of school or college	Town		
First name and initials	Surname		
Date of birth Day Month Year Candidate number	Number of seat		
All questions should be attempted. Necessary data will be found in the Chemistry Highe (1999 Edition). [This will be provided October/Nover The questions may be answered in any order but all ar	r and Advanced Higher Data Booklet nber 1998.] nswers are to be written in this answer		

Rough work, if any should be necessary, as well as the fair copy, is to be written in this book.

Rough work should be scored through when the fair copy has been written.

Additional space for answers and rough work will be found at the end of the book. If further space is required, supplementary sheets may be obtained from the invigilator and should be inserted inside the **front** cover of this booklet.

The size of the space provided for an answer should not be taken as an indication of how much to write. It is not necessary to use all the space.

Before leaving the examination room, you must give this book to the invigilator. If you do not, you may lose all the marks for this paper.







		Mauka	must not write in this margin
3.	Ammonia is now one of the world's most important chemicals, about two million tonnes being produced each year in the UK alone.	Marks	
	It is manufactured by the direct combination of nitrogen and hydrogen by the Haber Process.		
	$N_2(g) + 3H_2(g) \longrightarrow 2NH_3(g) \Delta H = -92 \text{ kJ mol}^{-1}$		
	The Haber Process is operated as a continuous process, and the reaction is never allowed to reach equilibrium.		
	(<i>a</i>) (i) What does the term "equilibrium" mean, when applied to a chemical reaction?		
	(ii) Why is the Haber Process not carried out at a very high temperature?	1	
		1	
	(<i>b</i>) State one advantage of operating the Haber Process as a continuous process.		
		1	
		(3)	

Candidate

Marks

4. The apparatus below was used by a student to find the enthalpies of combustion of alcohols.



(a) Write down the measurements the student should take.

(b) The enthalpies of combustion of three alcohols are shown in the table.

Alcohol	Enthalpy of combustion/kJ mol ⁻¹
methanol	-715
ethanol	-1371
propanol	-2010

Why is there a **regular** stepwise increase in the enthalpies of combustion from methanol to ethanol to propanol?

1 (3)

2



margin

Marks

6. The following triglyceride is found in some fats and oils.

$$\begin{array}{cccccc} H_{2}C - O - C - & (CH_{2})_{16}CH_{3} \\ & & \\ & & \\ & & \\ HC - O - C - & (CH_{2})_{7}CH &= & CH(CH_{2})_{7}CH_{3} \\ & & \\ & & \\ & & \\ H_{2}C - O - C - & (CH_{2})_{16}CH_{3} \\ & &$$

- (a) The hydrolysis of the triglyceride produces an alcohol and long chain fatty acids.
 - (i) Name the alcohol produced by the hydrolysis of the triglyceride.

(ii) Suggest why the sequence of fatty acids in the triglyceride can be referred to as S, O, S. (You may wish to refer to the data booklet.)

(b) What happens to triglyceride molecules in the conversion of oils to hardened fats?

1 (3)

1

1

Candidate must not write in this margin Marks 7. Urea is a substance found in human urine. The enzyme urease catalyses the hydrolysis of urea. urease enzyme CO₂ 2NH₃ $CO(NH_2)_2$ H₂O + + urea The concentration of urea in a sample can be estimated using an indicator as shown in the diagram. urea sample urease extract acidified gel +bromothymol blue indicator The bromothymol blue indicator is yellow below pH6 and blue above pH8.3. Draw the full structural formula for urea. *(a)* 1 *(b)* The initial yellow colour of the indicator changed to blue as the experiment proceeded. Explain fully the colours observed. 2



Candidate must not write in this margin Marks Esters are important compounds which have many applications. 8. Some of the instructions outlining the laboratory preparation of an ester are shown below. *(a)* Preparation of an Ester Mix 1 cm³ of the alkanol with 1 cm³ of the alkanoic acid in a test tube. 1 2 3 4 After 20 minutes, pour the contents of the test tube into a beaker containing sodium hydrogen carbonate solution. Add appropriate instructions for steps 2 and 3.

2

(b) The full structural formula for an ester is shown below.



Give the systematic name of the alkanol used in making this ester.

(c) State a use of esters.

1

Marks

Candidate must not write in this margin

9. In 1926, Hinshelwood and Green studied the reaction between nitric oxide and hydrogen at temperatures above 150 °C.

The equation for the reaction is:

2NO(g) + $2H_2(g)$ \rightarrow $N_2(g)$ + $2H_2O(g)$

A simplified diagram of their apparatus is shown below.



(a) Predict what will happen to the mercury levels as the reaction proceeds.

(b) The use of narrow glass tubing ensured that only a small volume of gas was outwith the reaction vessel.Suggest why this precaution was taken.

(c) Calculate the mass of nitrogen obtained under these conditions when 500 cm³ of nitric oxide reacted completely with hydrogen.
(Take the molar volume of nitrogen to be 25.0 litres mol⁻¹.)
(Show your working clearly.)

2

1

1

Marks

2

Candidate must not write in this margin

10. The boiling points of three halogens are shown in the table.

Halogen	Boiling point/°C
chlorine	-35
bromine	59
iodine	184

(a) Explain why the boiling points of the halogens increase down the group.



Marks

11. The enthalpy of lattice breaking for rubidium chloride is the enthalpy change for the following process.

RbCl(s) \longrightarrow $Rb^+(g)$ +

Cl (g)

This enthalpy change can be calculated using the enthalpy changes in the table below.

		Enthalpy change	Δ H/kJmol ⁻¹
1	RbCl(s)	\longrightarrow Rb ⁺ (aq) + Cl ⁻ (aq)	+17
2	Rb ⁺ (g)	\longrightarrow Rb ⁺ (aq)	-301
3	Cl ⁻ (g)	← Cl ⁻ (aq)	-364

(*a*) Name enthalpy change 1.

(b) Calculate the enthalpy of lattice breaking, in kJ mol⁻¹, for rubidium chloride.(Show your working clearly.)

2 (3)

1

12. Part of a workcard is shown.



A student carrying out this experiment passed the current through the solution for 10 minutes. *(b)* Calculate the mass of hydrogen produced. (Show your working clearly.)

3

Candidate must not write in this

Page fourteen

(5)

must not write in this margin

Marks

1

Candidate

13. The following reaction can be readily carried out in the laboratory.



(a) Why can this reaction be classified as oxidation?

(b) The alcohol does not react with bromine solution and the product does not react with Benedict's solution.Draw a structural formula for the product.

1 (2)



14.	(continue	d)	Marks	Candidate must not write in this margin
	(b) Sul	phuric acid is used in the manufacture of fertilisers.		
	(i) Write a balanced equation showing the formation of ammonium sulphate		
		from ammonia and sulphuric acid.		
	(i	i) Explain why ammonium sulphate dissolves in water to form an acidic solution.	1	

(5)

1

1 (2) must not write in this margin

15. Part of a polymer structure is shown below.



(a) Draw the structure for the monomer from which it is made.

(b) This polymer can be treated to make a polymer which conducts electricity. Why does it conduct electricity?

[C012/SQP006] 32

				Candidate must not write in this margin
16.	Addi Mark the d	tion of hydrogen chloride to an alkene can give two products. covnikoff observed that the hydrogen of the hydrogen chloride mainly attaches to the carbon atom of ouble bond which already has the most hydrogens directly attached to it.	Marks	
	(<i>a</i>)	Draw the full structural formula for the major product formed when hydrogen chloride reacts with propene.		
			1	
	(b)	Why is it not necessary to consider Markovnikoff's rule when hydrogen chloride reacts with but-2-ene?		
			1 (2)	
		Page nineteen		

Marks

1

17. The water in swimming pools can be kept sterile by the addition of chlorine which kills microorganisms. The chlorine levels in swimming pool water can be determined by titrating samples against acidified iron(II) sulphate solution. The reaction taking place is:

 $Cl_2(aq)$ + $2Fe^{2+}(aq)$ \longrightarrow $2Cl^{-}(aq)$ + $2Fe^{3+}(aq)$

(a) Write the ion-electron equation for the oxidation half reaction.

(b) A 100 cm³ sample of water from a swimming pool required 24.9 cm³ of 2.82×10^{-4} mol 1⁻¹ iron(II) sulphate solution to reach the end point. Calculate the chlorine concentration, in g 1⁻¹, in the swimming pool water. (Show your working clearly.)

[END OF QUESTION PAPER]



SPACE FOR ANSWERS

SPACE FOR ANSWERS