



National
Qualifications
SPECIMEN ONLY

S847/76/11

Mathematics
Paper 1 (Non-calculator)

Date — Not applicable

Duration — 1 hour 15 minutes



Total marks — 55

Attempt ALL questions.

You must NOT use a calculator.

To earn full marks you must show your working in your answers.

State the units for your answer where appropriate.

You will not earn marks for answers obtained by readings from scale drawings.

Write your answers clearly in the spaces provided in the answer booklet. The size of the space provided for an answer is not an indication of how much to write. You do not need to use all the space.

Additional space for answers is provided at the end of the answer booklet. If you use this space you must clearly identify the question number you are attempting.

Use blue or black ink.

Before leaving the examination room you must give your answer booklet to the Invigilator; if you do not, you may lose all the marks for this paper.



FORMULAE LIST

Circle

The equation $x^2 + y^2 + 2gx + 2fy + c = 0$ represents a circle centre $(-g, -f)$ and radius $\sqrt{g^2 + f^2 - c}$.

The equation $(x-a)^2 + (y-b)^2 = r^2$ represents a circle centre (a, b) and radius r .

Scalar product

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta, \text{ where } \theta \text{ is the angle between } \mathbf{a} \text{ and } \mathbf{b}$$

or $\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$ where $\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$.

Trigonometric formulae

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$= 2 \cos^2 A - 1$$

$$= 1 - 2 \sin^2 A$$

Table of standard derivatives

$f(x)$	$f'(x)$
$\sin ax$	$a \cos ax$
$\cos ax$	$-a \sin ax$

Table of standard integrals

$f(x)$	$\int f(x)dx$
$\sin ax$	$-\frac{1}{a} \cos ax + c$
$\cos ax$	$\frac{1}{a} \sin ax + c$

Total marks — 55

Attempt ALL questions


1. A curve has equation $y = x^2 - 4x + 7$.

Find the equation of the tangent to this curve at the point where $x = 5$.

4

2. A and B are the points $(-7, 3)$ and $(1, 5)$.

AB is a diameter of a circle.



Find the equation of this circle.

3

3. Line l_1 has equation $\sqrt{3}y - x = 0$.

(a) Line l_2 is perpendicular to l_1 . Find the gradient of l_2 .

2

(b) Calculate the angle l_2 makes with the positive direction of the x -axis.

2

4. The points A(0, 9, 7), B(5, $\overrightarrow{-1}$, 2), C(4, 1, 3) and D(x , -2, 2) are such that \vec{AB} is perpendicular to \vec{CD} .

Determine the value of x .

4

5. For the polynomial, $x^3 - 4x^2 + ax + b$

- $x - 1$ is a factor
- -12 is the remainder when it is divided by $x - 2$

(a) Determine the values of a and b .

5

(b) Hence solve $x^3 - 4x^2 + ax + b = 0$.

3

6. A sequence is generated by the recurrence relation $u_{n+1} = mu_n + 6$ where m is a constant.

(a) Given $u_1 = 28$ and $u_2 = 13$, find the value of m .

2

(b) (i) Explain why this sequence approaches a limit as $n \rightarrow \infty$.

1

(ii) Calculate this limit.

2


7. (a) Evaluate $\log_5 25$.

1

(b) Hence solve $\log_4 x + \log_4 (x - 6) = \log_5 25$, where $x > 6$.

5

8. Triangle ABD is right-angled at B with angles $BAC = p$ and $BAD = q$ and lengths as shown in the diagram below.



Show that the exact value of $\cos(q - p)$ is $\frac{19\sqrt{17}}{85}$.

5

9. The curve $y = f(x)$ is such that $\frac{dy}{dx} = 4x - 6x^2$. The curve passes through the point $(-1, 9)$. Express y in terms of x .

4

10. (a) Solve $\cos 2x^\circ - 3 \cos x^\circ + 2 = 0$ for $0 \leq x < 360$.

5

- (b) Hence solve $\cos 4x^\circ - 3 \cos 2x^\circ + 2 = 0$ for $0 \leq x < 360$.

2

[Turn over

11. Functions f and g are defined on the set of real numbers by $f(x) = x^3 - 1$ and $g(x) = 3x + 1$.

- (a) Find an expression for $k(x)$, where $k(x) = g(f(x))$. 2
- (b) If $h(k(x)) = x$, find an expression for $h(x)$. 3

[END OF SPECIMEN QUESTION PAPER]



National
Qualifications
SPECIMEN ONLY

S847/76/11

**Mathematics
Paper 1 (Non-calculator)**

Marking Instructions

These marking instructions have been provided to show how SQA would mark this specimen question paper.

The information in this publication may be reproduced to support SQA qualifications only on a non-commercial basis. If it is reproduced, SQA should be clearly acknowledged as the source. If it is to be used for any other purpose, written permission must be obtained from permissions@sqa.org.uk.

Where the publication includes materials from sources other than SQA (ie secondary copyright), this material should only be reproduced for the purposes of examination or assessment. If it needs to be reproduced for any other purpose it is the user's responsibility to obtain the necessary copyright clearance.

General marking principles for Higher Mathematics

Always apply these general principles. Use them in conjunction with the detailed marking instructions, which identify the key features required in candidates' responses.

For each question, the marking instructions are generally in two sections:

- *generic scheme – this indicates why each mark is awarded*
- *illustrative scheme – this covers methods which are commonly seen throughout the marking*

In general, you should use the illustrative scheme. Only use the generic scheme where a candidate has used a method not covered in the illustrative scheme.

- (a) Always use positive marking. This means candidates accumulate marks for the demonstration of relevant skills, knowledge and understanding; marks are not deducted for errors or omissions.
- (b) If a candidate response does not seem to be covered by either the principles or detailed marking instructions, and you are uncertain how to assess it, you must seek guidance from your team leader.
- (c) One mark is available for each •. There are no half marks.
- (d) If a candidate's response contains an error, all working subsequent to this error must still be marked. Only award marks if the level of difficulty in their working is similar to the level of difficulty in the illustrative scheme.
- (e) Only award full marks where the solution contains appropriate working. A correct answer with no working receives no mark, unless specifically mentioned in the marking instructions.
- (f) Candidates may use any mathematically correct method to answer questions, except in cases where a particular method is specified or excluded.
- (g) If an error is trivial, casual or insignificant, for example $6 \times 6 = 12$, candidates lose the opportunity to gain a mark, except for instances such as the second example in point (h) below.

- (h) If a candidate makes a transcription error (question paper to script or within script), they lose the opportunity to gain the next process mark, for example

This is a transcription error and so the mark is not awarded.

$$x^2 + 5x + 7 = 9x + 4$$

This is no longer a solution of a quadratic equation, so the mark is not awarded.

$$x - 4x + 3 = 0$$

$$x = 1$$

The following example is an exception to the above

This error is not treated as a transcription error, as the candidate deals with the intended quadratic equation. The candidate has been given the benefit of the doubt and all marks awarded.

$$x^2 + 5x + 7 = 9x + 4$$

$$x - 4x + 3 = 0$$

$$(x - 3)(x - 1) = 0$$

$$x = 1 \text{ or } 3$$

(i) **Horizontal/vertical marking**

If a question results in two pairs of solutions, apply the following technique, but only if indicated in the detailed marking instructions for the question.

Example:

$$\begin{array}{ll} \bullet^5 & \bullet^6 \\ \bullet^5 \quad x = 2 & x = -4 \\ \bullet^6 \quad y = 5 & y = -7 \end{array}$$

Horizontal: $\bullet^5 x = 2$ and $x = -4$
 $\bullet^6 y = 5$ and $y = -7$

Vertical: $\bullet^5 x = 2$ and $y = 5$
 $\bullet^6 x = -4$ and $y = -7$

You must choose whichever method benefits the candidate, **not** a combination of both.

- (j) In final answers, candidates should simplify numerical values as far as possible unless specifically mentioned in the detailed marking instruction. For example

$\frac{15}{12}$ must be simplified to $\frac{5}{4}$ or $1\frac{1}{4}$ $\frac{43}{1}$ must be simplified to 43

$\frac{15}{0.3}$ must be simplified to 50 $\frac{4}{5}$ must be simplified to $\frac{4}{15}$

$\sqrt{64}$ must be simplified to 8*

*The square root of perfect squares up to and including 144 must be known.

- (k) Do not penalise candidates for any of the following, unless specifically mentioned in the detailed marking instructions:
- working subsequent to a correct answer
 - correct working in the wrong part of a question
 - legitimate variations in numerical answers/algebraic expressions, for example angles in degrees rounded to nearest degree
 - omission of units
 - bad form (bad form only becomes bad form if subsequent working is correct), for example

$(x^3 + 2x^2 + 3x + 2)(2x + 1)$ written as

$$(x^3 + 2x^2 + 3x + 2) \times 2x + 1$$

$$= 2x^4 + 5x^3 + 8x^2 + 7x + 2$$

gains full credit

- repeated error within a question, but not between questions or papers

- (l) In any ‘Show that...’ question, where candidates have to arrive at a required result, the last mark is not awarded as a follow-through from a previous error, unless specified in the detailed marking instructions.
- (m) You must check all working carefully, even where a fundamental misunderstanding is apparent early in a candidate’s response. You may still be able to award marks later in the question so you must refer continually to the marking instructions. The appearance of the correct answer does not necessarily indicate that you can award all the available marks to a candidate.
- (n) You should mark legible scored-out working that has not been replaced. However, if the scored-out working has been replaced, you must only mark the replacement working.
- (o) If candidates make multiple attempts using the same strategy and do not identify their final answer, mark all attempts and award the lowest mark. If candidates try different valid strategies, apply the above rule to attempts within each strategy and then award the highest mark.

For example:

Strategy 1 attempt 1 is worth 3 marks.	Strategy 2 attempt 1 is worth 1 mark.
Strategy 1 attempt 2 is worth 4 marks.	Strategy 2 attempt 2 is worth 5 marks.
From the attempts using strategy 1, the resultant mark would be 3.	From the attempts using strategy 2, the resultant mark would be 1.

In this case, award 3 marks.

Marking instructions for each question

Question			Generic scheme	Illustrative scheme	Max mark
1.			<ul style="list-style-type: none"> •¹ differentiate •² calculate gradient •³ find the value of y •⁴ find equation of tangent 	<ul style="list-style-type: none"> •¹ $2x - 4$ •² 6 •³ 12 •⁴ $y = 6x - 18$ 	4
2.			<ul style="list-style-type: none"> •¹ find the centre •² calculate the radius •³ state equation of circle 	<ul style="list-style-type: none"> •¹ $(-3, 4)$ •² $\sqrt{17}$ •³ $(x+3)^2 + (y-4)^2 = 17$ or equivalent 	3
3.	(a)		<ul style="list-style-type: none"> •¹ find gradient l_1 •² state gradient l_2 	<ul style="list-style-type: none"> •¹ $\frac{1}{\sqrt{3}}$ •² $-\sqrt{3}$ 	2
	(b)		<ul style="list-style-type: none"> •³ using $m = \tan \theta$ •⁴ calculating angle 	<ul style="list-style-type: none"> •³ $\tan \theta = -\sqrt{3}$ •⁴ $\theta = \frac{2\pi}{3}$ or 120° 	2

Question		Generic scheme	Illustrative scheme	Max mark
4.		<ul style="list-style-type: none"> •¹ find \vec{CD} •² find \vec{AB} •³ equate scalar product to zero •⁴ calculate value of x 	<ul style="list-style-type: none"> •¹ $\begin{pmatrix} x-4 \\ -3 \\ -1 \end{pmatrix}$ •² $\begin{pmatrix} 5 \\ -10 \\ -5 \end{pmatrix}$ •³ $5(x-4) + (-10)(-3) + (-5)(-1) = 0$ •⁴ $x = -3$ 	4

Question		Generic scheme	Illustrative scheme	Max mark	
5.	(a)	<ul style="list-style-type: none"> •¹ use appropriate strategy •² obtain an expression for a and b •³ obtain a second expression for a and b •⁴ find the value of a or b •⁵ find the second value 	<ul style="list-style-type: none"> •¹ $(1)^3 - 4(1)^2 + a(1) + b = 0$ •² $a + b = 3$ •³ $2a + b = -4$ •⁴ $a = -7$ or $b = 10$ •⁵ $b = 10$ or $a = -7$ 	5	
	(b)	<ul style="list-style-type: none"> •⁶ obtain quadratic factor •⁷ complete factorisation •⁸ state solutions 	<ul style="list-style-type: none"> •⁶ $(x^2 - 3x - 10)$ •⁷ $(x-1)(x-5)(x+2)$ •⁸ $x = 1, x = 5, x = -2$ 	3	
6.	(a)	<ul style="list-style-type: none"> •¹ interpret information •² solve to find m 	<ul style="list-style-type: none"> •¹ $13 = 28m + 6$ •² $m = \frac{1}{4}$ 	2	
	(b)	(i)	<ul style="list-style-type: none"> •³ state condition 	<ul style="list-style-type: none"> •³ a limit exists as $-1 < \frac{1}{4} < 1$ 	1
		(ii)	<ul style="list-style-type: none"> •⁴ know how to calculate limit •⁵ calculate limit 	<ul style="list-style-type: none"> •⁴ $L = \frac{1}{4}L + 6$ •⁵ $L = 8$ 	2

Question		Generic scheme	Illustrative scheme	Max mark
7.	(a)	• ¹ state value	• ¹ 2	1
	(b)	<ul style="list-style-type: none"> •¹ use laws of logarithms •² link to part (a) •³ use laws of logarithms •⁴ write in standard quadratic form •⁵ solve for x and identify appropriate solution 	<ul style="list-style-type: none"> •¹ $\log_4 x(x-6)$ •² $\log_4 x(x-6) = 2$ •³ $x(x-6) = 4^2$ •⁴ $x^2 - 6x - 16 = 0$ •⁵ 8 	5
8.		<ul style="list-style-type: none"> •¹ calculate lengths AC and AD •² select appropriate formula and express in terms of p and q •³ calculate two of $\cos p, \cos q, \sin p, \sin q$ •⁴ calculate other two and substitute into formula •⁵ arrange into required form 	<ul style="list-style-type: none"> •¹ $AC = \sqrt{17}$ and $AD = 5$ stated or implied by •³ •² $\cos q \cos p + \sin q \sin p$ stated or implied by •⁴ •³ $\cos p = \frac{4}{\sqrt{17}}$, $\cos q = \frac{4}{5}$ $\sin p = \frac{1}{\sqrt{17}}$, $\sin q = \frac{3}{5}$ •⁴ $\frac{4}{5} \times \frac{4}{\sqrt{17}} + \frac{3}{5} \times \frac{1}{\sqrt{17}}$ •⁵ $\frac{19}{5\sqrt{17}} \times \frac{\sqrt{17}}{\sqrt{17}} = \frac{19\sqrt{17}}{85}$ or $\frac{19}{5\sqrt{17}} = \frac{19\sqrt{17}}{5 \times 17} = \frac{19\sqrt{17}}{85}$ 	5

Question		Generic scheme	Illustrative scheme	Max mark
9.		<ul style="list-style-type: none"> •¹ know to and start to integrate •² complete integration •³ substitute for x and y •⁴ state expression for y 	<ul style="list-style-type: none"> •¹ eg $y = \frac{4}{2}x^2 \dots$ •² $y = \frac{4}{2}x^2 - \frac{6}{3}x^3 + c$ •³ $9 = 2(-1)^2 - 2(-1)^3 + c$ •⁴ $y = 2x^2 - 2x^3 + 5$ 	4
10.	(a)	<ul style="list-style-type: none"> •¹ use double angle formula •² express as a quadratic in $\cos x^\circ$ •³ start to solve •⁴ reduce to equations in $\cos x^\circ$ only •⁵ process solutions 	<p>Method 1: Using factorisation</p> <ul style="list-style-type: none"> •¹ $2\cos^2 x^\circ - 1 \dots$ stated or implied by •² •² $2\cos^2 x^\circ - 3\cos x^\circ + 1 = 0$ •³ $(2\cos x^\circ - 1)(\cos x^\circ - 1)$ <p>Method 2: Using quadratic formula</p> <ul style="list-style-type: none"> •¹ $2\cos^2 x^\circ - 1 \dots$ stated or implied by •² •² $2\cos^2 x^\circ - 3\cos x^\circ + 1 = 0$ stated explicitly •³ $\frac{-(-3) \pm \sqrt{(-3)^2 - 4 \times 2 \times 1}}{2 \times 2}$ <p>In both methods:</p> <ul style="list-style-type: none"> •⁴ $\cos x^\circ = \frac{1}{2}$ and $\cos x^\circ = 1$ •⁵ $0, 60, 300$ or •⁴ $\cos x = 1$ and $x = 0$ •⁵ $\cos x^\circ = \frac{1}{2}$ and $x = 60$ or 300 	5
	(b)	<ul style="list-style-type: none"> •⁶ interpret relationship with (a) •⁷ state valid values 	<ul style="list-style-type: none"> •⁶ $2x = 0$ and 60 and 300 •⁷ $0, 30, 150, 180, 210$ and 330 	2

Question		Generic scheme	Illustrative scheme	Max mark
11.	(a)	<ul style="list-style-type: none"> •¹ interpret notation •² complete process 	<ul style="list-style-type: none"> •¹ $g(x^3 - 1)$ •² $3x^3 - 2$ 	2
	(b)	<ul style="list-style-type: none"> •³ start to rearrange for x •⁴ rearrange •⁵ state expression for $h(x)$ 	<ul style="list-style-type: none"> •³ $3x^3 = y + 2$ •⁴ $x = \sqrt[3]{\frac{y+2}{3}}$ •⁵ $h(x) = \sqrt[3]{\frac{x+2}{3}}$ 	3

[END OF SPECIMEN MARKING INSTRUCTIONS]