



National  
Qualifications  
SPECIMEN ONLY

**S847/76/12**

**Mathematics  
Paper 2**

Date — Not applicable

Duration — 1 hour 30 minutes

---

**Total marks — 65**

Attempt ALL questions.

**You may use a calculator.**

To earn full marks you must show your working in your answers.

State the units for your answer where appropriate.

You will not earn marks for answers obtained by readings from scale drawings.

Write your answers clearly in the spaces provided in the answer booklet. The size of the space provided for an answer is not an indication of how much to write. You do not need to use all the space.

Additional space for answers is provided at the end of the answer booklet. If you use this space you must clearly identify the question number you are attempting.

Use **blue** or **black** ink.

Before leaving the examination room you must give your answer booklet to the Invigilator; if you do not, you may lose all the marks for this paper.



## FORMULAE LIST

### Circle

The equation  $x^2 + y^2 + 2gx + 2fy + c = 0$  represents a circle centre  $(-g, -f)$  and radius  $\sqrt{g^2 + f^2 - c}$ .

The equation  $(x - a)^2 + (y - b)^2 = r^2$  represents a circle centre  $(a, b)$  and radius  $r$ .

### Scalar product

$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta$ , where  $\theta$  is the angle between  $\mathbf{a}$  and  $\mathbf{b}$

or  $\mathbf{a} \cdot \mathbf{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$  where  $\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$ .

### Trigonometric formulae

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$= 2 \cos^2 A - 1$$

$$= 1 - 2 \sin^2 A$$

### Table of standard derivatives

$f(x)$	$f'(x)$
$\sin ax$	$a \cos ax$
$\cos ax$	$-a \sin ax$


### Table of standard integrals

$f(x)$	$\int f(x) dx$
$\sin ax$	$-\frac{1}{a} \cos ax + c$
$\cos ax$	$\frac{1}{a} \sin ax + c$

Total marks — 65  
Attempt ALL questions

1. The vertices of triangle ABC are  $A(-5, 7)$ ,  $B(-1, -5)$  and  $C(13, 3)$  as shown in the diagram.

The broken line represents the altitude from C.




- |     |   |   |
|-----|---|---|
| (a) | Find the equation of the altitude from C.   | 3 |
| (b) | Find the equation of the median from B.   | 3 |
| (c) | Find the coordinates of the point of intersection of the altitude from C and the median from B. | 2 |
- 
- |    |  |   |
|----|--|---|
| 2. | Find $\int \frac{4x^3 + 1}{x^2} dx, x \neq 0.$ | 4 |
|----|--|---|

[Turn over

3. The diagram shows the curve with equation  $y = f(x)$ , where  $f(x) = kx(x+a)(x+b)$ .


The curve passes through  $(-1, 0)$ ,  $(0, 0)$ ,  $(1, 2)$  and  $(2, 0)$ .



Find the values of  $a$ ,  $b$  and  $k$ .

3

4. D,OABC is a square-based pyramid as shown.



- O is the origin and  $OA = 4$  units.
- M is the mid-point of OA.
- $\vec{OD} = 2\mathbf{i} + 2\mathbf{j} + 6\mathbf{k}$


(a) Express  $\vec{DB}$  and  $\vec{DM}$  in component form.

3

(b) Find the size of angle BDM.

5

5. The line with equation  $y = 2x + 3$  is a tangent to the curve with equation  $y = x^3 + 3x^2 + 2x + 3$  at  $A(0, 3)$ , as shown.



The line meets the curve again at  $B(-3, -3)$ .


Find the area enclosed by the line and the curve.

5

6. (a) Express  $3x^2 + 24x + 50$  in the form  $a(x+b)^2 + c$ . 3
- (b) Given that  $f(x) = x^3 + 12x^2 + 50x - 11$ , find  $f'(x)$ . 2
- (c) Hence, or otherwise, explain why the curve with equation  $y = f(x)$  is strictly increasing for all values of  $x$ . 2

[Turn over

7. The diagram below shows the graph of a quartic  $y = h(x)$ , with stationary points at  $(0, 5)$  and  $(2, 2)$ .



On separate diagrams sketch the graphs of:

(a)  $y = 2 - h(x)$ .


2

(b)  $y = h'(x)$ .

3

8. A design for a new grain container is in the shape of a cylinder with a hemispherical roof and a flat circular base. The radius of the cylinder is  $r$  metres, and the height is  $h$  metres.

The volume of the **cylindrical** part of the container needs to be 100 cubic metres.



- (a) Given that the curved surface area of a hemisphere of radius  $r$  is  $2\pi r^2$  show that the surface area of metal needed to build the grain container is given by:

$$A = \frac{200}{r} + 3\pi r^2 \text{ square metres}$$

3

- (b) Determine the value of  $r$  which minimises the amount of metal needed to build the container.

6

9. Given that

$$\int_{\frac{\pi}{8}}^a \sin\left(4x - \frac{\pi}{2}\right) dx = \frac{1}{2}, \quad 0 \leq a < \frac{\pi}{2},$$

calculate the value of  $a$ .

6

[Turn over


10. Show that  $\frac{\sin 2x}{2 \cos x} - \sin x \cos^2 x = \sin^3 x$ , where  $0 < x < \frac{\pi}{2}$ .

3

11. (a) Show that the points  $A(-7, -2)$ ,  $B(2, 1)$  and  $C(17, 6)$  are collinear.

3

Three circles with centres  $A$ ,  $B$  and  $C$  are drawn inside a circle with centre  $D$  as shown.



The circles with centres  $A$ ,  $B$  and  $C$  have radii  $r_A$ ,  $r_B$  and  $r_C$  respectively.

- $r_A = \sqrt{10}$
- $r_B = 2r_A$
- $r_C = r_A + r_B$

(b) Determine the equation of the circle with centre  $D$ .

4

[END OF SPECIMEN QUESTION PAPER]





National  
Qualifications  
SPECIMEN ONLY

**S847/76/12**

**Mathematics  
Paper 2**

## Marking Instructions

---

These marking instructions have been provided to show how SQA would mark this specimen question paper.

The information in this publication may be reproduced to support SQA qualifications only on a non-commercial basis. If it is reproduced, SQA should be clearly acknowledged as the source. If it is to be used for any other purpose, written permission must be obtained from [permissions@sqa.org.uk](mailto:permissions@sqa.org.uk).

Where the publication includes materials from sources other than SQA (ie secondary copyright), this material should only be reproduced for the purposes of examination or assessment. If it needs to be reproduced for any other purpose it is the user's responsibility to obtain the necessary copyright clearance.

## General marking principles for Higher Mathematics

*Always apply these general principles. Use them in conjunction with the detailed marking instructions, which identify the key features required in candidates' responses.*

*For each question, the marking instructions are generally in two sections:*

- *generic scheme – this indicates why each mark is awarded*
- *illustrative scheme – this covers methods which are commonly seen throughout the marking*

*In general, you should use the illustrative scheme. Only use the generic scheme where a candidate has used a method not covered in the illustrative scheme.*

- (a) Always use positive marking. This means candidates accumulate marks for the demonstration of relevant skills, knowledge and understanding; marks are not deducted for errors or omissions.
- (b) If a candidate response does not seem to be covered by either the principles or detailed marking instructions, and you are uncertain how to assess it, you must seek guidance from your team leader.
- (c) One mark is available for each •. There are no half marks.
- (d) If a candidate's response contains an error, all working subsequent to this error must still be marked. Only award marks if the level of difficulty in their working is similar to the level of difficulty in the illustrative scheme.
- (e) Only award full marks where the solution contains appropriate working. A correct answer with no working receives no mark, unless specifically mentioned in the marking instructions.
- (f) Candidates may use any mathematically correct method to answer questions, except in cases where a particular method is specified or excluded.
- (g) If an error is trivial, casual or insignificant, for example  $6 \times 6 = 12$ , candidates lose the opportunity to gain a mark, except for instances such as the second example in point (h) below.

- (h) If a candidate makes a transcription error (question paper to script or within script), they lose the opportunity to gain the next process mark, for example

This is a transcription error and so the mark is not awarded.

This is no longer a solution of a quadratic equation, so the mark is not awarded.

$$x^2 + 5x + 7 = 9x + 4$$

$$x - 4x + 3 = 0$$

$$x = 1$$

The following example is an exception to the above

This error is not treated as a transcription error, as the candidate deals with the intended quadratic equation. The candidate has been given the benefit of the doubt and all marks awarded.

$$x^2 + 5x + 7 = 9x + 4$$

$$x - 4x + 3 = 0$$

$$(x - 3)(x - 1) = 0$$

$$x = 1 \text{ or } 3$$

- (i) **Horizontal/vertical marking**

If a question results in two pairs of solutions, apply the following technique, but only if indicated in the detailed marking instructions for the question.

Example:

$$\begin{array}{cc} \bullet^5 & \bullet^6 \\ \bullet^5 & x = 2 \quad x = -4 \\ \bullet^6 & y = 5 \quad y = -7 \end{array}$$

Horizontal:  $\bullet^5 x = 2 \text{ and } x = -4$   
 $\bullet^6 y = 5 \text{ and } y = -7$

Vertical:  $\bullet^5 x = 2 \text{ and } y = 5$   
 $\bullet^6 x = -4 \text{ and } y = -7$

You must choose whichever method benefits the candidate, **not** a combination of both.

- (j) In final answers, candidates should simplify numerical values as far as possible unless specifically mentioned in the detailed marking instruction. For example

$\frac{15}{12}$  must be simplified to  $\frac{5}{4}$  or  $1\frac{1}{4}$        $\frac{43}{1}$  must be simplified to 43

$\frac{15}{0.3}$  must be simplified to 50       $\frac{4\cancel{5}}{3}$  must be simplified to  $\frac{4}{15}$

$\sqrt{64}$  must be simplified to 8\*

\*The square root of perfect squares up to and including 144 must be known.

(k) Do not penalise candidates for any of the following, unless specifically mentioned in the detailed marking instructions:

- working subsequent to a correct answer
- correct working in the wrong part of a question
- legitimate variations in numerical answers/algebraic expressions, for example angles in degrees rounded to nearest degree
- omission of units
- bad form (bad form only becomes bad form if subsequent working is correct), for example

$(x^3 + 2x^2 + 3x + 2)(2x + 1)$  written as

$(x^3 + 2x^2 + 3x + 2) \times 2x + 1$

$= 2x^4 + 5x^3 + 8x^2 + 7x + 2$

gains full credit

- repeated error within a question, but not between questions or papers

(l) In any 'Show that...' question, where candidates have to arrive at a required result, the last mark is not awarded as a follow-through from a previous error, unless specified in the detailed marking instructions.

(m) You must check all working carefully, even where a fundamental misunderstanding is apparent early in a candidate's response. You may still be able to award marks later in the question so you must refer continually to the marking instructions. The appearance of the correct answer does not necessarily indicate that you can award all the available marks to a candidate.

(n) You should mark legible scored-out working that has not been replaced. However, if the scored-out working has been replaced, you must only mark the replacement working.

(o) If candidates make multiple attempts using the same strategy and do not identify their final answer, mark all attempts and award the lowest mark. If candidates try different valid strategies, apply the above rule to attempts within each strategy and then award the highest mark.

For example:

Strategy 1 attempt 1 is worth 3 marks.	Strategy 2 attempt 1 is worth 1 mark.
Strategy 1 attempt 2 is worth 4 marks.	Strategy 2 attempt 2 is worth 5 marks.
From the attempts using strategy 1, the resultant mark would be 3.	From the attempts using strategy 2, the resultant mark would be 1.

In this case, award 3 marks.

### Marking instructions for each question



Question			Generic scheme	Illustrative scheme	Max mark
1.	(a)		<ul style="list-style-type: none"> <li>•<sup>1</sup> calculate gradient of AB</li> <li>•<sup>2</sup> use property of perpendicular lines</li> <li>•<sup>3</sup> determine equation of altitude</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> <math>m_{AB} = -3</math></li> <li>•<sup>2</sup> <math>m_{alt} = \frac{1}{3}</math></li> <li>•<sup>3</sup> <math>x - 3y = 4</math></li> </ul>	3
	(b)		<ul style="list-style-type: none"> <li>•<sup>4</sup> calculate midpoint of AC</li> <li>•<sup>5</sup> calculate gradient of median</li> <li>•<sup>6</sup> determine equation of median</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>4</sup> (4,5)</li> <li>•<sup>5</sup> <math>m_{BM} = 2</math></li> <li>•<sup>6</sup> <math>y = 2x - 3</math></li> </ul>	3
	(c)		<ul style="list-style-type: none"> <li>•<sup>7</sup> find <math>x</math> or <math>y</math> coordinate</li> <li>•<sup>8</sup> find remaining coordinate</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>7</sup> <math>x = 1</math> or <math>y = -1</math></li> <li>•<sup>8</sup> <math>y = -1</math> or <math>x = 1</math></li> </ul>	2
2.			<ul style="list-style-type: none"> <li>•<sup>1</sup> write in integrable form</li> <li>•<sup>2</sup> integrate one term</li> <li>•<sup>3</sup> integrate other term</li> <li>•<sup>4</sup> complete integration and simplify</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> <math>4x + x^{-2}</math></li> <li>•<sup>2</sup> eg <math>\frac{4}{2}x^2 + \dots</math></li> <li>•<sup>3</sup> <math>\dots \frac{x^{-1}}{-1}</math></li> <li>•<sup>4</sup> <math>2x^2 - x^{-1} + c</math></li> </ul>	4
3.			<ul style="list-style-type: none"> <li>•<sup>1</sup> value of <math>a</math></li> <li>•<sup>2</sup> value of <math>b</math></li> <li>•<sup>3</sup> calculate <math>k</math></li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> 1</li> <li>•<sup>2</sup> -2</li> <li>•<sup>3</sup> -1</li> </ul>	3

Question			Generic scheme	Illustrative scheme	Max mark
4.	(a)		<ul style="list-style-type: none"> <li>•<sup>1</sup> state components of <math>\overrightarrow{DB}</math></li> <li>•<sup>2</sup> state coordinates of M</li> <li>•<sup>3</sup> state components of <math>\overrightarrow{DM}</math></li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> <math>\begin{pmatrix} 2 \\ 2 \\ -6 \end{pmatrix}</math></li> <li>•<sup>2</sup> (2,0,0) stated or implied by •<sup>3</sup></li> <li>•<sup>3</sup> <math>\begin{pmatrix} 0 \\ -2 \\ -6 \end{pmatrix}</math></li> </ul>	3
	(b)		<ul style="list-style-type: none"> <li>•<sup>4</sup> evaluate <math>\overrightarrow{DB} \cdot \overrightarrow{DM}</math></li> <li>•<sup>5</sup> evaluate <math> \overrightarrow{DB} </math></li> <li>•<sup>6</sup> evaluate <math> \overrightarrow{DM} </math></li> <li>•<sup>7</sup> use scalar product</li> <li>•<sup>8</sup> calculate angle</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>4</sup> 32</li> <li>•<sup>5</sup> <math>\sqrt{44}</math></li> <li>•<sup>6</sup> <math>\sqrt{40}</math></li> <li>•<sup>7</sup> <math>\cos BDM = \frac{32}{\sqrt{44}\sqrt{40}}</math></li> <li>•<sup>8</sup> 40.3° or 0.703 rads</li> </ul>	5

Question			Generic scheme	Illustrative scheme	Max mark
5.			<ul style="list-style-type: none"> <li>•<sup>1</sup> know to integrate and interpret limits</li> <li>•<sup>2</sup> use 'upper – lower'</li> <li>•<sup>3</sup> integrate</li> <li>•<sup>4</sup> substitute limits</li> <li>•<sup>5</sup> evaluate area</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> <math>\int_{-3}^0 \dots dx</math></li> <li>•<sup>2</sup> <math>\int_{-3}^0 (x^3 + 3x^2 + 2x + 3) - (2x + 3) dx</math></li> <li>•<sup>3</sup> <math>\frac{1}{4}x^4 + x^3</math></li> <li>•<sup>4</sup> <math>0 - \left( \frac{1}{4}(-3)^4 + (-3)^3 \right)</math></li> <li>•<sup>5</sup> <math>\frac{27}{4}</math> units<sup>2</sup></li> </ul>	5

Question			Generic scheme	Illustrative scheme	Max mark
6.	(a)		<b>Method 1</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> identify common factor</li> <li>•<sup>2</sup> complete the square</li> <li>•<sup>3</sup> process for <math>c</math> and write in required form</li> </ul>	<b>Method 1</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> <math>3(x^2 + 8x \dots)</math> stated or implied by •<sup>2</sup></li> <li>•<sup>2</sup> <math>3(x+4)^2 \dots</math></li> <li>•<sup>3</sup> <math>3(x+4)^2 + 2</math></li> </ul>	3
			<b>Method 2</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> expand completed square form</li> <li>•<sup>2</sup> equate coefficients</li> <li>•<sup>3</sup> process for <math>b</math> and <math>c</math> and write in required form</li> </ul>	<b>Method 2</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> <math>ax^2 + 2abx + ab^2 + c</math></li> <li>•<sup>2</sup> <math>a = 3, 2ab = 24, ab^2 + c = 50</math></li> <li>•<sup>3</sup> <math>3(x+4)^2 + 2</math></li> </ul>	3
	(b)		<ul style="list-style-type: none"> <li>•<sup>4</sup> differentiate two terms</li> <li>•<sup>5</sup> complete differentiation</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>4</sup> <math>3x^2 + 24x \dots</math></li> <li>•<sup>5</sup> <math>\dots + 50</math></li> </ul>	2
	(c)		<b>Method 1</b> <ul style="list-style-type: none"> <li>•<sup>6</sup> link with (a) and identify sign of <math>(x+4)^2</math></li> <li>•<sup>7</sup> communicate reason</li> </ul>	<b>Method 1</b> <ul style="list-style-type: none"> <li>•<sup>6</sup> <math>f'(x) = 3(x+4)^2 + 2</math> and <math>(x+4)^2 \geq 0 \forall x</math></li> <li>•<sup>7</sup> <math>\therefore 3(x+4)^2 + 2 &gt; 0 \Rightarrow</math> always strictly increasing</li> </ul>	2
			<b>Method 2</b> <ul style="list-style-type: none"> <li>•<sup>6</sup> identify minimum value of <math>f'(x)</math></li> <li>•<sup>7</sup> communicate reason</li> </ul>	<b>Method 2</b> <ul style="list-style-type: none"> <li>•<sup>6</sup> eg minimum value = 2 or annotated sketch</li> <li>•<sup>7</sup> <math>2 &gt; 0 \therefore (f'(x) &gt; 0) \Rightarrow</math> always strictly increasing</li> </ul>	2



Question			Generic scheme	Illustrative scheme	Max mark
7.	(a)		<ul style="list-style-type: none"> <li>•<sup>1</sup> evidence of reflecting in <math>x</math>-axis</li> <li>•<sup>2</sup> vertical translation of 2 units identifiable from graph</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> reflection of graph in <math>x</math>-axis</li> <li>•<sup>2</sup> graph moves parallel to <math>y</math>-axis by 2 units upwards</li> </ul> 	2
	(b)		<ul style="list-style-type: none"> <li>•<sup>3</sup> identify roots</li> <li>•<sup>4</sup> interpret point of inflexion</li> <li>•<sup>5</sup> complete cubic curve</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>3</sup> 0 and 2 only</li> <li>•<sup>4</sup> turning point at (2, 0)</li> <li>•<sup>5</sup> cubic passing through origin with negative gradient</li> </ul> 	3

Question			Generic scheme	Illustrative scheme	Max mark
8.	(a)		<ul style="list-style-type: none"> <li>•<sup>1</sup> equate volume to 100</li> <li>•<sup>2</sup> obtain an expression for <math>h</math></li> <li>•<sup>3</sup> demonstrate result</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> <math>V = \pi r^2 h = 100</math></li> <li>•<sup>2</sup> <math>h = \frac{100}{\pi r^2}</math></li> <li>•<sup>3</sup> <math>A = \pi r^2 + 2\pi r^2 + 2\pi r \times \frac{100}{\pi r^2}</math> leading to  <math>A = \frac{200}{r} + 3\pi r^2</math></li> </ul>	3
	(b)		<ul style="list-style-type: none"> <li>•<sup>4</sup> start to differentiate</li> <li>•<sup>5</sup> complete differentiation</li> <li>•<sup>6</sup> set derivative to zero</li> <li>•<sup>7</sup> obtain <math>r</math></li> <li>•<sup>8</sup> verify nature of stationary point</li> <li>•<sup>9</sup> interpret and communicate result</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>4</sup> <math>A'(r) = 6\pi r \dots</math></li> <li>•<sup>5</sup> <math>A'(r) = 6\pi r - \frac{200}{r^2}</math></li> <li>•<sup>6</sup> <math>6\pi r - \frac{200}{r^2} = 0</math></li> <li>•<sup>7</sup> <math>r = \sqrt[3]{\frac{100}{3\pi}} (\approx 2.20) \text{ metres}</math></li> <li>•<sup>8</sup> table of signs for a derivative when <math>r = 2.1974 \dots</math></li> <li>•<sup>9</sup> minimum when <math>r \approx 2.20 \text{ (m)}</math></li> </ul> <p style="text-align: center;">or</p> <ul style="list-style-type: none"> <li>•<sup>8</sup> <math>A''(r) = 6\pi + \frac{400}{r^3}</math></li> <li>•<sup>9</sup> <math>A''(2.1974 \dots) &gt; 0 \therefore \text{minimum when } r \approx 2.20 \text{ (m)}</math></li> </ul>	6

Question			Generic scheme	Illustrative scheme	Max mark
9.			<ul style="list-style-type: none"> <li>•<sup>1</sup> start to integrate</li> <li>•<sup>2</sup> complete integration</li> <li>•<sup>3</sup> process limits</li> <li>•<sup>4</sup> simplify numeric term and equate to <math>\frac{1}{2}</math></li> <li>•<sup>5</sup> start to solve equation</li> <li>•<sup>6</sup> solve for <math>a</math></li> </ul>	<ul style="list-style-type: none"> <li>•<sup>1</sup> <math>-\frac{1}{4}\cos\ldots</math></li> <li>•<sup>2</sup> <math>-\frac{1}{4}\cos\left(4x-\frac{\pi}{2}\right)</math></li> <li>•<sup>3</sup> <math>-\frac{1}{4}\cos\left(4a-\frac{\pi}{2}\right)+\frac{1}{4}\cos\left(\frac{4\pi}{8}-\frac{\pi}{2}\right)</math></li> <li>•<sup>4</sup> <math>-\frac{1}{4}\cos\left(4a-\frac{\pi}{2}\right)+\frac{1}{4}=\frac{1}{2}</math></li> <li>•<sup>5</sup> <math>\cos\left(4a-\frac{\pi}{2}\right)=-1</math></li> <li>•<sup>6</sup> <math>a=\frac{3\pi}{8}</math></li> </ul>	6
10.			<b>Method 1</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> substitute for <math>\sin 2x</math></li> <li>•<sup>2</sup> simplify and factorise</li> <li>•<sup>3</sup> substitute for <math>1-\cos^2 x</math> and simplify</li> </ul>	<b>Method 1</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> <math>\frac{2\sin x \cos x}{2\cos x} - \sin x \cos^2 x</math> stated explicitly as above or in a simplified form of the above</li> <li>•<sup>2</sup> <math>\sin x(1-\cos^2 x)</math></li> <li>•<sup>3</sup> <math>\sin x \times \sin^2 x</math> leading to <math>\sin^3 x</math></li> </ul>	3
			<b>Method 2</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> substitute for <math>\sin 2x</math></li> <li>•<sup>2</sup> simplify and substitute for <math>\cos^2 x</math></li> <li>•<sup>3</sup> expand and simplify</li> </ul>	<b>Method 2</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> <math>\frac{2\sin x \cos x}{2\cos x} - \sin x \cos^2 x</math> stated explicitly as above or in a simplified form of the above</li> <li>•<sup>2</sup> <math>\sin x - \sin x(1-\sin^2 x)</math></li> <li>•<sup>3</sup> <math>\sin x - \sin x + \sin^3 x</math> leading to <math>\sin^3 x</math></li> </ul>	3

Question			Generic scheme	Illustrative scheme	Max mark
11.	(a)		<b>Method 1</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> calculate <math>m_{AB}</math></li> <li>•<sup>2</sup> calculate <math>m_{BC}</math></li> <li>•<sup>3</sup> interpret result and state conclusion</li> </ul>	<b>Method 1</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> eg <math>m_{AB} = \frac{3}{9} = \frac{1}{3}</math></li> <li>•<sup>2</sup> eg <math>m_{BC} = \frac{5}{15} = \frac{1}{3}</math></li> <li>•<sup>3</sup> ... <math>\Rightarrow</math> AB and BC are parallel (common direction), B is a common point, hence A, B and C are collinear.</li> </ul>	3
			<b>Method 2</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> calculate an appropriate vector, eg <math>\overrightarrow{AB}</math></li> <li>•<sup>2</sup> calculate a second vector, eg <math>\overrightarrow{BC}</math> and compare</li> <li>•<sup>3</sup> interpret result and state conclusion</li> </ul>	<b>Method 2</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> eg <math>\overrightarrow{AB} = \begin{pmatrix} 9 \\ 3 \end{pmatrix}</math></li> <li>•<sup>2</sup> eg <math>\overrightarrow{BC} = \begin{pmatrix} 15 \\ 5 \end{pmatrix} \therefore \overrightarrow{AB} = \frac{3}{5}\overrightarrow{BC}</math></li> <li>•<sup>3</sup> ... <math>\Rightarrow</math> AB and BC are parallel (common direction), B is a common point, hence A, B and C are collinear.</li> </ul>	3
			<b>Method 3</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> calculate <math>m_{AB}</math></li> <li>•<sup>2</sup> find equation of line and substitute point</li> <li>•<sup>3</sup> communication</li> </ul>	<b>Method 3</b> <ul style="list-style-type: none"> <li>•<sup>1</sup> <math>m_{AB} = \frac{3}{9} = \frac{1}{3}</math></li> <li>•<sup>2</sup> eg, <math>y - 1 = \frac{1}{3}(x - 2)</math> leading to <math>6 - 1 = \frac{1}{3}(17 - 2)</math></li> <li>•<sup>3</sup> since C lies on line A, B and C are collinear</li> </ul>	3
	(b)		<ul style="list-style-type: none"> <li>•<sup>4</sup> find radius</li> <li>•<sup>5</sup> determine an appropriate ratio</li> <li>•<sup>6</sup> find centre</li> <li>•<sup>7</sup> state equation of circle</li> </ul>	<ul style="list-style-type: none"> <li>•<sup>4</sup> <math>6\sqrt{10}</math></li> <li>•<sup>5</sup> eg 2 : 3 or <math>\frac{2}{5}</math> (using B and C) or 3 : 5 or <math>\frac{8}{5}</math> (using A and C)</li> <li>•<sup>6</sup> (8,3)</li> <li>•<sup>7</sup> <math>(x - 8)^2 + (y - 3)^2 = 360</math></li> </ul>	4

[END OF SPECIMEN MARKING INSTRUCTIONS]